

## For the Period June 6 to 12, 2017

Seeding has essentially wrapped up in the province, according to Saskatchewan Agriculture's Weekly Crop Report. Producers have 99 per cent of the crop in the ground, ahead of the five-year (2012-2016) seeding average of 97 per cent for this time of year. The northwest region is the furthest behind with 97 per cent seeded. Across the province, there are still a few fields of oats and barley, as well as some greenfeed and silage, left to be seeded.

The much-needed rainfall received this week will help replenish topsoil moisture and allow crops to advance. While much of the north has surplus topsoil moisture at this time, many areas in the south were relieved to see rain as crops, hay land and pastures were in need of significant moisture.

Prior to the recent rainfall, topsoil moisture conditions were deteriorating in southern and central areas of the province. Provincially, cropland topsoil moisture is currently rated as seven per cent surplus, 52 per cent adequate, 35 per cent short and six per cent very short. Hay land and pasture topsoil moisture is rated as four per cent surplus, 43 per cent adequate, 41 per cent short and 12 per cent very short.

Overall crop development has improved with the warm and wet weather. Sixty per cent of fall cereals, 62 per cent of spring cereals, 53 of oilseeds and 70 per cent of pulse crops are at their normal stages of development for this time of year. The majority of crops are in fair-to-excellent condition.

Persistent wind has damaged crops and delayed in-crop weed control operations in many areas. Cutworms, flea beetles, localized flooding, hail and lack of moisture have also caused damage this past week.

Provincial Crop Development			
	% Ahead	% Normal	% Behind
Fall Cereals	22	60	18
Spring Cereals	5	62	33
Oilseeds	3	53	44
Pulse Crops	5	70	25

Producers are busy completing seeding, controlling weeds and getting ready for haying.

SaskPower reports that there have been 149 reports of power line contact this year, with 19 incidents reported in June.

### One year ago

Producers had seeded 99.5 per cent of the 2016 crop. Crops were advancing well and much needed rain was received across the province.

Follow the 2017 Crop Report on Twitter @SKAgriculture

### Seeding Progress in SK Per cent seeded All Crops

June 12, 2017	99
June 13, 2016	100
June 15, 2015	100
June 16, 2014	95
June 10, 2013	96
June 11, 2012	96
5 year avg. (2012-2016)	97
10 year avg. (2007-2016)	95

For further information, contact Shannon Friesen, PAg,  
Cropping Management Specialist, Moose Jaw, Regional Services Branch,  
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: [cropreport@gov.sk.ca](mailto:cropreport@gov.sk.ca).  
Also available on the Ministry of Agriculture website at [www.saskatchewan.ca/crop-report](http://www.saskatchewan.ca/crop-report).



### **Southeastern Saskatchewan:**

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville, Minton and Lake Alma areas

Seeding has wrapped up in the region, although there are still some greenfeed and silage crops left to be seeded. In-crop weed control is underway in areas with active weed growth; however, strong winds have been delaying producers for several weeks now.

<b>Southeast Crop Development</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	24	60	16
Spring Cereals	14	52	34
Oilseeds	9	43	48
Pulse Crops	13	42	45

Rainfall was welcomed in many areas over the past week as topsoil moisture conditions were rapidly deteriorating. Rainfall this past week ranged from small amounts to 41 mm in the Maryfield area. The Carnduff area reported 4 mm of rain, the Kisbey area 26 mm, the Moosomin and Glenavon areas 12 mm, the Fillmore area 7 mm, the Weyburn area 3 mm, the Indian Head area 10 mm and the Moose Jaw area 20 mm. The Moose Jaw area has reported receiving the most precipitation (96 mm) in the region since April 1.

Despite the rainfall a week ago, topsoil moisture conditions have not significantly improved; however, significant rainfall over the past couple of days may help improve conditions. Cropland topsoil moisture is rated as one per cent surplus, 49 per cent adequate, 33 per cent short and 17 per cent very short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 40 per cent adequate, 37 per cent short and 22 per cent very short. Crop District 2A is reporting that 56 per cent of cropland and 71 per cent of hay land and pasture are very short topsoil moisture at this time, and CD 3ASE is reporting that 75 per cent of cropland and 80 per cent of hay land and pasture are very short topsoil moisture.

Overall, the majority of crops are in fair-to-good condition, although many crops in drier areas are suffering from lack of moisture. Many crops are quite short and the recent rainfall may not help crop development or yield. There are reports of thin and stunted winter cereal crops and hay fields that are prematurely heading-out. Even with the moisture, hay fields in many areas will yield less than could be expected as the rain came too late. Some producers are expecting the haying season to be shorter than normal this year.

The majority of crop damage this past week was due to lack of moisture, wind and insects such as cutworms and flea beetles in the canola crops. Gophers and alfalfa weevils have also caused some issues in the area.

Farmers are busy trying to control weeds, picking rocks and getting ready for haying.

### Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Seeding has all but wrapped up for producers in the southwest, but there are still some greenfeed and silage crops to be seeded in the coming weeks. In-crop weed control continues in many areas; however, strong winds have been delaying producers for several weeks now.

Southwest Crop Development			
	% Ahead	% Normal	% Behind
Fall Cereals	19	60	21
Spring Cereals	3	74	23
Oilseeds	0	72	28
Pulse Crops	4	79	17

Much of the region received small amounts of rain last week, although the Cabri area reported 12 mm. The Fife Lake and Climax areas received 2 mm of rain last week, the Limerick area 9 mm, the Mossbank area 5 mm, the Mortlach, Eastend and Blumenhof areas 6 mm, the Hazenmore area 10 mm, the Webb and Maple Creek areas 7 mm and the Leader area 8 mm. The Tyner area has reported receiving the most precipitation (83 mm) in the region since April 1.

Some areas in the region are much drier than others and will need a significant rain soon to help crops emerge and grow. The rainfall over the past couple of days may help improve topsoil moisture conditions in the region which have rapidly deteriorated over the past few weeks due to lack of moisture and strong winds. Cropland topsoil moisture is rated as 39 per cent adequate, 54 per cent short and seven per cent very short. Hay land and pasture topsoil moisture is rated as 25 per cent adequate, 60 per cent short and 15 per cent very short. Crop District 3BS is reporting that 24 per cent of cropland and 36 per cent of hay land and pasture are very short topsoil moisture at this time.

Overall, the majority of crops are in fair-to-good condition, although many crops in drier areas are suffering from lack of moisture. Many crops are quite short and the recent rainfall may not help crop development or yield. Livestock producers have indicated that some hay fields are stunted and prematurely heading-out. Yields will likely be compromised as the recent rain may have been too late for some areas. Some producers are expecting the haying season to be shorter than normal this year.

The majority of crop damage this past week was due to lack of moisture, wind and insects such as cutworms and flea beetles in the canola crops. Gophers have also caused some issues in the region.

Farmers are busy trying to control weeds, picking rocks and getting ready for haying.

**East-Central Saskatchewan:**

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Seeding has essentially wrapped up in the region and producers are now busy with in-crop herbicide applications when the winds are calm. Some oat, barley and greenfeed crops remain to be seeded in the coming weeks.

Rainfall last week ranged from trace amounts to 40 mm in the Rose Valley area. The

Esterhazy area reported 8 mm of rain, the Yorkton and Fife Lake areas 6 mm, the Kelliher area 2 mm, the Quill Lake area 24 mm, the Lumsden area 16 mm, the Earl Grey area 4 mm, the Craik area 11 mm and the Chamberlain area 15 mm. The Rose Valley area has reported receiving the most precipitation (94 mm) in the region since April 1.

<b>East-Central Crop Development</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	11	79	10
Spring Cereals	3	65	32
Oilseeds	2	59	39
Pulse Crops	5	74	21

Despite this rainfall, strong winds have been drying out the topsoil and many later-seeded crops will need a good rain in order to establish themselves. The rainfall in the last couple of days should help with plant growth, although there are concerns that significant rainfall may flood some already-wet fields. At this time, cropland topsoil moisture is rated as three per cent surplus, 50 per cent adequate, 44 per cent short and three per cent very short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 40 per cent adequate, 51 per cent short and eight per cent very short. Crop District 5B is reporting that 10 per cent of cropland currently has surplus topsoil moisture.

Overall, the majority of crops are in fair-to-good condition, although crop development has been slower than normal. Many crops and hay fields in drier areas are suffering from lack of moisture and are very short in stature. Even with the recent moisture, hay fields in many areas are expected to yield less than could be expected.

Along with lack of moisture and strong winds, frost, hail and insects such as flea beetles in canola have caused most of the crop damage this past week. Gophers and alfalfa weevils have also caused some issues in the area.

Farmers are busy trying to control weeds and getting ready for haying.

**West-Central Saskatchewan:**

- Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B – Kerrobert, Macklin, Wilkie and Biggar areas

Producers in the west-central region have essentially completed seeding and are actively applying in-crop herbicides when weather permits. There will be some greenfeed crops such as oats and barley seeded in the coming weeks.

<b>West-Central Crop Development</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	19	75	6
Spring Cereals	1	70	29
Oilseeds	1	65	34
Pulse Crops	1	75	24

Rainfall in the region ranged from trace amounts to 30 mm in the Macklin area. The Eyebrow, Unity and Tramping Lake areas reported 10 mm of rain last week, the Rosthern area 6 mm, the Rosetown area 7 mm, the Marengo area 13 mm and the Kindersley area 16 mm. The Battleford area has reported receiving the most precipitation (137 mm) in the region since April 1.

Although recent rainfall has helped to replenish topsoil moisture in some areas of the region, strong winds have been drying out fields at the same time. More rain is needed to help crops emerge and establish and for hay fields and pastures to green-up. Cropland topsoil moisture is rated as one per cent surplus, 62 per cent adequate, 36 per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 55 per cent adequate, 37 per cent short and seven per cent very short.

The majority of crops in the region are in fair-to-excellent condition, although crop development in some areas has been slower than normal. Later-seeded crops in drier areas are suffering from lack of moisture and are very short in stature. Livestock producers are concerned that hay yields will be lower than expected if additional rainfall is not received soon.

Most crop damage this past week is attributed to lack of moisture, localized flooding, strong winds and insects such as cutworms and flea beetles. Producers continue to spray for these insects and to re-seed patches in fields.

Farmers are busy controlling weeds and insects, hauling grain and picking rocks.

**Northeastern Saskatchewan:**

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

Other than some oat and barley fields that still need to be seeded, producers in much of the region are essentially done. However, some areas near Nipawin and Carrot River are dealing with excess moisture which will result in some fields remaining unseeded. Localized flooding has caused road access issues and has damaged farmyards.

<b>Northeast Crop Development</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	8	42	50
Spring Cereals	3	42	55
Oilseeds	2	37	61
Pulse Crops	5	40	55

Rainfall in the region ranged from small amounts to 63 mm in the Nipawin area. The Tisdale area reported 14 mm, the Vonda area 1 mm, the Melfort area 15 mm, the Birch Hills area 5 mm, the Garrick area 21 mm and the Spruce Home area 32 mm. The Nipawin area has reported receiving the most precipitation (266 mm) in the region since April 1.

Topsoil moisture conditions have deteriorated this past week due to the heavy rainfall. Cropland topsoil moisture is rated as 37 per cent surplus, 56 per cent adequate, four per cent short and three per cent very short. Hay land and pasture topsoil moisture is rated as 26 per cent surplus, 69 per cent adequate, four per cent short and one per cent very short. Crop districts 8A and 9AE are reporting that 63 per cent and 43 per cent, respectively, of cropland have surplus topsoil moisture at this time.

Crops in the region are in fair-to-excellent condition, although crop development has been slower than normal due to excess moisture and delayed seeding. Producers were able to combine many of the unharvested fields from 2016, although some fields remain unharvested. In-crop spraying operations are now underway as weather and time permit.

The majority of crop damage this week is attributed to localized flooding, frost, wind and insects such as flea beetles.

Farmers are busy trying to complete seeding, controlling weeds and getting ready for haying.

**Northwestern Saskatchewan:**

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Producers in the northwestern region are nearly ready wrap up seeding, as they now have 97 per cent of the crop in the ground. If the weather co-operates, many of the remaining fields should be seeded within the coming week. However, some fields are expected to remain unseeded due to excess moisture. In-crop spraying operations are also underway as weather and time permit.

<b>Northwest Crop Development</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	45	55	0
Spring Cereals	4	50	46
Oilseeds	3	49	48
Pulse Crops	4	73	23

Large amounts of rain were reported last week, with the Meadow Lake area reporting a range of 44 to 48 mm. There were also reports of some areas receiving up to four inches of rain in a short period of time. The Duck Lake area reported 22 mm, the Rabbit Lake area 35 mm, the Debden area 32 mm, the Neilburg area 28 mm and the Turtleford area 20 mm. The Pierceland area has reported receiving the most precipitation (237 mm) in the region since April 1.

Despite the rain, topsoil moisture conditions have improved overall in the region. Cropland topsoil moisture is rated as 20 per cent surplus, 68 per cent adequate and 12 per cent short. Hay land and pasture topsoil moisture is rated as 17 per cent surplus, 72 per cent adequate and 11 per cent short. Crop District 9B is reporting that 33 per cent of cropland and 26 per cent of hay land and pasture have surplus topsoil moisture at this time.

Overall, crops are in fair-to-excellent condition. Crop development has been slower than normal but is quickly catching up thanks to warmer weather. Rainfall will be needed in many areas in the coming weeks to help crops establish and for hay land and pasture to grow. Even with the recent moisture, hay fields in many areas will yield less than could be expected.

Localized flooding has caused issues in many far northwestern areas, and hail has damaged some emerging crops. Strong winds have caused some damage, along with insects such as flea beetles.

Farmers are busy trying to complete seeding, controlling weeds and getting ready for haying.



### Crop Conditions - June 12, 2017

Provincial							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	10	8	16	17	11	11	12
% good	66	66	59	42	64	62	53
% fair	16	19	22	32	19	22	27
% poor	6	6	3	8	5	2	7
% very poor	2	1	0	1	1	3	1
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	3	11	2	6	19	18	0
% good	45	53	46	54	58	51	54
% fair	39	27	41	30	18	27	33
% poor	10	7	9	8	4	4	13
% very poor	3	2	2	2	1	0	0
Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	10	4	8	0	4	6	8
% good	63	73	58	29	64	51	39
% fair	17	14	27	45	24	33	35
% poor	8	8	6	18	6	8	16
% very poor	2	1	1	8	2	2	2
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	2	5	2	7	10	5	0
% good	32	38	43	46	59	43	20
% fair	45	37	37	34	23	41	70
% poor	14	13	16	10	7	10	10
% very poor	7	7	2	3	1	1	0
Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	17	0	19	23	6	6	20
% good	62	41	45	41	35	61	50
% fair	13	57	30	29	54	31	20
% poor	8	2	6	7	5	2	10
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	2	11	0	0	20	19	0
% good	39	52	41	100	50	53	53
% fair	40	26	46	0	23	24	33
% poor	19	10	10	0	7	4	14
% very poor	0	1	3	0	0	0	0
East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	8	10	6	4	9	5	12
% good	77	83	68	87	66	67	73
% fair	15	7	25	9	20	27	15
% poor	0	0	1	0	5	1	0
% very poor	0	0	0	0	0	0	0

East-central (continued)							
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	3	4	0	5	10	6	0
% good	54	60	77	80	77	78	85
% fair	41	30	23	15	13	15	15
% poor	2	6	0	0	0	1	0
% very poor	0	0	0	0	0	0	0
West-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	20	40	19	15	16	16	19
% good	58	33	59	55	69	62	41
% fair	22	19	21	29	14	21	38
% poor	0	8	1	1	1	1	2
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	0	17	10	0	25	28	0
% good	68	59	61	80	58	47	100
% fair	31	19	28	20	16	24	0
% poor	1	5	1	0	1	1	0
% very poor	0	0	0	0	0	0	0
Northeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	21	28	14	21	0
% good	0	10	51	57	62	56	94
% fair	0	20	18	12	13	9	5
% poor	0	13	8	3	8	2	1
% very poor	100	57	2	0	3	12	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	6	14	0	24	19	15	N/A
% good	85	49	72	66	52	60	N/A
% fair	8	25	20	9	13	20	N/A
% poor	1	9	8	1	1	5	N/A
% very poor	0	3	0	0	15	0	N/A
Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	72	34	32	N/A	17	17	0
% good	28	60	56	N/A	71	73	100
% fair	0	6	12	N/A	12	10	0
% poor	0	0	0	N/A	0	0	0
% very poor	0	0	0	N/A	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	20	23	N/A	N/A	23	45	N/A
% good	60	60	N/A	N/A	70	43	N/A
% fair	20	16	N/A	N/A	7	12	N/A
% poor	0	1	N/A	N/A	0	0	N/A
% very poor	0	0	N/A	N/A	0	0	N/A

# Crop Districts and Rural Municipalities in Saskatchewan

- Regional Service Office
- Regional Satellite Office
- Crop District
- Rural Municipality




Data Source:  
Crop Districts - Saskatchewan Ministry of Agriculture

Geomatic Services, Ministry of Agriculture June 10, 2014

# Weekly Rainfall

from June 6 to June 12, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

# Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period June 6 to 12, 2017


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	4	59	4A	49	White Valley	6	55	7A	287	St. Andrews	7	63
	3	Enniskillen	11	63		51	Reno	3.29	44.66		288	Pleasant Valley	9	65
	31	Storhokoaks	N/A	10		79	Arlington	NIL	51		290 A	Kindersley	4	66
	32	Reciprocity	11	78		109	Carmichael	N/A	78		290 B	Kindersley	N/A	40
	33	Moose Creek	N/A	55		110	Piapot	7	61.5		290 C	Kindersley	N/A	24
	34	Browning	9	56		111	Maple Creek	N/A	49		292	Milton	13	86
	61	Antler	N/A	5	4B	139 A	Gull Lake	N/A	37		317 A	Marriott	N/A	N/A
	64	Brock	26	74		139 B	Gull Lake	NIL	35.5		317 B	Marriott	3	64
	65	Tecumseh	N/A	39		142	Enterprise	N/A	N/A		318	Mountain View	4	106
1B	91	Maryfield	41	70		169	Pittville	N/A	34		320 A	Oakdale	16	64.5
	94	Hazelwood	24.03	24.03		231	Happyland	8	71		320 B	Oakdale	10	68
	122	Martin	12	30	5A	183	Fertile Belt	8	41		321	Prairiedale	11	82
	123	Silverwood	9	37		211 A	Churchbridge	NIL	53	7B	347	Biggar	NIL	80
	124	Kingsley	11	53		211 B	Churchbridge	4	39		350 A	Mariposa	10.14	105.64
	125 A	Chester	7	39		211 C	Churchbridge	N/A	NIL		350 B	Mariposa	4	81
	125 B	Chester	12	45		213	Saltcoats	N/A	42		351	Progress	6	117
	151	Rocanville	7	38		218	Cupar	N/A	N/A		352	Heart's Hill	6	106
	154 A	Elcapo	4	27		241	Calder	3	33		377	Glenside	NIL	82
	154 B	Elcapo	9	31		243	Wallace	NIL	13		378	Rosemount	NIL	109
	155	Wolseley	7	31.25		244	Orkney	6	18		379	Reford	N/A	86
2A	67	Weyburn	3	24		245 A	Garry	NIL	34		381	Grass Lake	N/A	64.5
	68	Brokenshell	4	29		245 B	Garry	4	53		382	Eye Hill	30	105
	96	Fillmore	6.5	10.25		245 C	Garry	2	53.5		409 A	Buffalo	1	137
	97	Wellington	8	21		246 A	Ituna Bon Accord	1	40		409 B	Buffalo	3	119
2B	127 A	Francis	8.5	50.5		246 B	Ituna Bon Accord	2.5	48.5		410	Round Valley	10	126.18
	127 B	Francis	7.5	28.75		247	Kellross	2	51	8A	395	Porcupine	N/A	87
	129	Bratt's Lake	4	23		248	Touchwood	N/A	36.5		397	Barrier Valley	13.69	81.45
	131 A	Baildon	4	63		271	Cote	NIL	65		428	Star City	9	74
	131 B	Baildon	12	96	5B	273	Sliding Hills	NIL	48		456	Arborfield	18	125
	156 A	Indian Head	10.19	39.02		277	Emerald	6	81		457	Connaught	18	140
	156 B	Indian Head	10	65		305	Invermay	NIL	70		486	Moose Range	40	147
	159	Sherwood	11	37		307	Elfros	NIL	45		487	Nipawin	63	265.5
	160 A	Pense	10	26		308 A	Big Quill	NIL	29	8B	369	St. Peter	15	54
	160 B	Pense	10	32.5		308 B	Big Quill	5	44		370 A	Humboldt	3	57
	161	Moose Jaw	20	50		331	Livingston	3	77		370 B	Humboldt	15	45
	162	Caron	11	38.5		335	Hazel Dell	NIL	26		371	Bayne	1	62
	191	Marquis	8	44		336	Sasman	2	49		372	Grant	1	89.37
3ASE	38 A	Laurier	0.4	16.9		337	Lakeview	N/A	46.5		400	Three Lakes	8	55
	38 B	Laurier	7	23		338	Lakeside	24	73		429 A	Flett's Springs	10	81
	39	The Gap	14	24		366	Kelvington	22	80		429 B	Flett's Springs	15	92.5
3ASW	10	Happy Valley	2	48		367	Ponass Lake	40	93.5		459	Kinistino	5	60
	12	Poplar Valley	2	49	6A	368	Spalding	18	70		460	Birch Hills	5.29	67.17
	40	Bengough	N/A	N/A		190 A	Dufferin	N/A	45	9AE	488	Torch River	21	114
	42	Willow Bunch	8	47		190 B	Dufferin	10	58		491	Buckland	32	55.5
	43	Old Post	NIL	39		190 C	Dufferin	16	69		520	Paddockwood	3	130
	73 A	Stonehenge	9	57.89		190 D	Dufferin	NIL	47	9AW	521	Lakeland	3	130
	73 B	Stonehenge	N/A	64		219 A	Longlaketon	4	44		406	Mayfield	0	59
3AN	102	Lake Johnston	4.59	44.32		219 B	Longlaketon	10	68		435	Redberry	N/A	120
	103	Sutton	6	47		220	McKillop	N/A	45		436	Douglas	NIL	97
	132 A	Hillsborough	6	31.5		221 A	Sarnia	13.39	66.74		463	Duck Lake	22	105.62
	132 B	Hillsborough	8	82		221 B	Sarnia	15.23	50.85		466	Meeting Lake	18	159
	193	Eyebrow	9	35		222	Craik	8	58		467 A	Round Hill	12	125
	17	Val Marie	N/A	20.97		251	Big Arm	N/A	34.5		467 B	Round Hill	35	129
3BS	18	Lone Tree	2	29		252	Arm River	11	51		467 C	Round Hill	27	162
	75	Pinto Creek	10	58		279	Mount Hope	2.29	37.78		494	Canwood	32	132
	76	Auvergne	5	55		282	McCraney	7	63	9B	438	Battle River	2	126
	77	Wise Creek	NIL	46		312	Morris	NIL	54.5		440	Hillsdale	28	140
	78	Grassy Creek	7	55.48		313	Lost River	N/A	48		442	Manitou Lake	13	151.12
	105	Glenbain	2	40		339	Leroy	0.8	38.98		498 A	Parkdale	25	166
	106	Whiska Creek	7	53		340	Wolverine	8	69		498 B	Parkdale	18	113.5
	107	Lac Pelletier	6	44		341	Viscount	NIL	55		499 A	Mervin	20	161
	108	Bone Creek	6	58		343 A	Blucher	NIL	35		501 A	Frenchman Butte	17	201
	138 A	Webb	7	49		223 A	Huron	10	60		501 B	Frenchman Butte	15	129
3BN	138 B	Webb	4	27	6B	223 B	Huron	N/A	46		501 C	Frenchman Butte	21	199
	165	Morse	N/A	67.5		284 A	Rudy	9	89		502	Britannia	24.5	197.5
	166	Excelsior	7	53		284 B	Rudy	NIL	57.5		561	Loon Lake	N/A	167
	168 A	Riverside	8	45		284 C	Rudy	N/A	47		588 A	Meadow Lake	48	195
	168 B	Riverside	11.75	45.44		285	Fertile Valley	4	57		588 B	Meadow Lake	44	222
	226	Victory	N/A	30		286	Milden	N/A	61		622	Beaver River	25	236.7
	228 A	Lacadena	8	83		314	Dundurn	NIL	57					
	228 B	Lacadena	N/A	6.5		344 A	Corman Park	NIL	77					
	257	Monet	6.5	65		344 B	Corman Park	N/A	NIL					
						346	Perdue	NIL	57					
						376	Eagle Creek	NIL	87					
						403	Rosthern	6	92					

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Municipality No: A, B, C and D - more than one reporter

# Cumulative Rainfall


from April 1 to June 12, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

# Cropland Topsoil Moisture Conditions


June 12, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

# Hay and Pasture Topsoil Moisture Conditions

## June 12, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

