

## For the Period June 20 to 26, 2017

Despite the cool and dry conditions experienced so far, the majority of crops are in good to excellent condition, according to Saskatchewan Agriculture's Weekly Crop Report.

Sixty-five per cent of winter wheat, 54 per cent of fall rye, 70 per cent of spring wheat, 49 per cent of durum, 68 per cent of oats, 64 per cent of barley, 60 per cent of canola, 50 per cent of flax, 63 per cent of lentils, 61 per cent of soybean, 65 per cent of peas, 34 per cent of chickpeas and 26 per cent of mustard are in good to excellent condition. Approximately 87 per cent of fall cereals, 62 per cent of spring cereals, 55 per cent of oilseed crops and 67 per cent of pulses are at or ahead of their normal stage of development for this time of year.

Precipitation across the province varied widely last week, ranging from negligible amounts to 57 mm in the Nipawin area. Provincially, topsoil moisture on cropland is rated as six per cent surplus, 61 per cent adequate, 28 per cent short and five per cent very short. Topsoil moisture on hay land and pasture is rated as five per cent surplus, 53 per cent adequate, 32 per cent short and 10 per cent very short. Many areas, particularly in the south, are in need of rain to help crops, hay and pasture grow.

There were reports of frost in some areas of the south, and producers are assessing the damage.

Haying is underway. Four per cent of the hay crop is cut and two per cent is baled or ensiled. Haying is furthest advanced in the southwest where 11 per cent of the crop has been cut. Hay quality is rated as six per cent excellent, 50 per cent good, 34 per cent fair and 10 per cent poor. Quality is lowest in the southeast where 21 per cent of the hay is rated as poor, thanks to less-than-ideal growing conditions during most of the spring.

Crop damage this week has been attributed to flooding, frost, wind and insects such as cutworms and alfalfa weevils. There are also reports of gopher damage to crops. Producers have nearly finished herbicide applications, but progress has been slowed by windy conditions.

Producers are scouting for disease, applying fungicides and cutting hay.

SaskPower has received 29 reports of farm equipment coming in contact with power lines so far in June. There have been 168 incidents in 2017.

For further information, contact Shannon Friesen, PAg,  
Cropping Management Specialist, Moose Jaw, Regional Services Branch,  
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: [cropreport@gov.sk.ca](mailto:cropreport@gov.sk.ca).  
Also available on the Ministry of Agriculture website at [www.saskatchewan.ca/crop-report](http://www.saskatchewan.ca/crop-report).

### One year ago

Crops were nicely advancing and were in good-to-excellent condition overall. Haying was underway with seven per cent of the hay crop cut and two per cent baled.

Follow the 2017 Crop Report on Twitter  
[@SKAgriculture](https://twitter.com/SKAgriculture)

### Provincial Crop Development - June 26, 2017

	% Ahead	% Normal	% Behind
Fall Cereals	13	74	13
Spring Cereals	4	58	38
Oilseeds	3	52	45
Pulse Crops	5	62	33

**Southeastern Saskatchewan:**

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu’Appelle areas
- Crop District 3ASE – Radville, Minton and Lake Alma areas

<b>Southeast Crop Development - June 26, 2017</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	16	72	12
Spring Cereals	10	57	33
Oilseeds	8	47	45
Pulse Crops	9	62	29

In-crop weed control is underway and is nearing completion in some areas after several weeks of delay due to strong winds and wet conditions.

Haying operations are underway in the region. Three per cent of the hay crop has been cut and two per cent baled or ensiled. Hay quality is rated as four per cent excellent, 38 per cent good, 37 per cent fair and 21 per cent poor. Due to less-than-ideal growing conditions, hay and pasture yields are expected to be less than normal.

Rainfall ranged from small amounts to 12 mm in the Briercrest area. The Glenavon area, which received a small amount of rain this week, holds the record (157 mm) for the greatest amount of precipitation in the region since April 1.

Topsoil moisture condition on cropland is rated as three per cent surplus, 61 per cent adequate, 32 per cent short, and four per cent very short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 57 per cent adequate, 33 per cent short, and nine per cent very short. Crop districts 3ASE (Ceylon-Radville area) and 2A (Weyburn area) report that 57 and 65 per cent of cropland, respectively, is short to very short topsoil moisture, and 90 and 77 per cent of hay land and pasture, respectively, are short to very short topsoil moisture.

Overall, most crops are in good-to-excellent condition and development is normal or ahead of schedule for this time of year. However, due to unfavourable growing conditions, 45 per cent of the oilseeds, 29 per cent of the pulse crops, 33 per cent of spring cereals, and 12 per cent of fall cereals are reported to be behind their normal development stage for this time of the year. Producers are hoping for calmer days so they can complete spraying and warm days to help invigorate crops, hay and pasture.

Crop damage this past week was attributed to flooding, wind and insects, mainly cutworms.

**Southwestern Saskatchewan:**

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current , Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Strong winds and dry conditions have delayed crop, hay and pasture development in the region. Haying operations are underway, and 10 per cent of the crop has been cut and five per cent baled or ensiled. Hay quality is rated as four per cent excellent, 46 per cent good, 42 per cent fair and eight per cent poor. Due to cool and dry growing conditions, hay and pasture yields are expected to be less than normal.

<b>Southwest Crop Development - June 26, 2017</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	3	87	10
Spring Cereals	5	60	35
Oilseeds	4	54	42
Pulse Crops	5	51	44

Rainfall across the region ranged from small amounts to 16 mm in the Swift Current area. The Moose Jaw area, which received 6 mm of rainfall this past week, holds the record (120 mm) for the greatest amount of precipitation in the regions since April 1.

Topsoil moisture on cropland is rated as 29 per cent adequate, 57 per cent short and 14 per cent very short. Hay land and pasture topsoil moisture is rated as 20 per cent adequate, 53 per cent short, and 27 per cent very short. Crop districts 3ASW (Rockglen area) and 3BS (Shaunavon area) report that 97 and 86 per cent of cropland, respectively, is short to very short topsoil moisture, and 97 and 94 per cent of hay land and pasture, respectively, are short to very short topsoil moisture. Crop District 4B (Maple Creek-Consul area) reports that 75 per cent of cropland is short to very short topsoil moisture and 100 per cent of hay land and pasture are short topsoil moisture.

While most crops are in good-to-excellent condition and development is normal or ahead of schedule for this time of year, the unfavourable growing conditions have slowed down development in some areas. Forty-two per cent of the oilseeds, 44 per cent of the pulse crops, 35 per cent of spring cereals and 10 per cent of fall cereals are reported to be behind their normal stage of development for this time of the year. Producers are hoping for calmer and warm days to help invigorate crops, hay and pasture.

Crop damage this past week was attributed to dry conditions, wind and frost.

**East-Central Saskatchewan:**

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Strong winds and dry and cool conditions have delayed crop, hay and pasture development in the region. Haying operations are underway and two per cent of the crop has been cut and one per cent baled or ensiled. Hay quality is rated as 50 per cent good, 35 per cent fair and 15 per cent poor. Due to cool and dry growing conditions, hay and pasture yields are expected to be less than normal. Some areas are reporting poor crop development as a result of cool and wet conditions.

<b>East-Central Crop Development - June 26, 2017</b>			
	% Ahead	% Normal	% Behind
Fall Cereals	12	67	21
Spring Cereals	3	56	41
Oilseeds	3	51	46
Pulse Crops	3	69	28

Rainfall across the region ranged from small amounts to 17 mm in the Kelvington area. The Kelvington area also holds the record (137 mm) for the greatest amount of precipitation in the region since April 1.

Topsoil moisture on cropland is rated as four per cent surplus, 75 per cent adequate, 18 per cent short and three per cent very short. Hay land and pasture topsoil moisture is rated as two per cent surplus, 70 per cent adequate, 24 per cent short and four per cent very short. Crop District 6A (Allan area) reports that 40 per cent of cropland and 51 per cent of hay land and pasture are short to very short topsoil moisture.

The unfavourable growing conditions have slowed crop development in some areas, with 46 per cent of oilseeds, 28 per cent of pulses, 41 per cent of spring cereals and 21 per cent of fall cereals reported to be behind their normal stages of development for this time of the year. Most producers are hoping for calm and warm days as well as rainfall to help invigorate crops, hay and pasture.

Crop damage this past week was attributed to dry conditions, wind, flooding, frost and flea beetles.

### West-Central Saskatchewan:

- Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B – Kerrobert, Macklin, Wilkie and Biggar areas

Strong winds and dry and cool conditions have delayed crop, hay and pasture development in the region. Haying operations are underway and three per cent of the crop has been cut. Hay quality is rated as 10 per cent excellent, 58 per cent good and 32 per cent fair. Due to cool and dry growing conditions, hay and pasture yields are expected to be less than normal.

West-Central Crop Development - June 26, 2017			
	% Ahead	% Normal	% Behind
Fall Cereals	15	79	6
Spring Cereals	1	69	30
Oilseeds	1	69	30
Pulse Crops	1	77	22

Rainfall across the region ranged from small amounts to 25 mm in the Kindersley area. The Battleford area holds the record (155 mm) for the greatest amount of precipitation in the region since April 1.

Topsoil moisture on cropland is rated as one per cent surplus, 71 per cent adequate, 26 per cent short and two per cent very short. Hay land and pasture topsoil moisture is rated as 62 per cent adequate, 33 per cent short and five per cent very short. Crop District 6B (Outlook area) reports that 43 per cent of cropland and 49 per cent of hay land and pasture are short to very short topsoil moisture.

The unfavourable growing conditions have slowed crop development in some areas, with 30 per cent of oilseeds, 22 per cent of pulses, 30 per cent of spring cereals and six per cent of fall cereals reported to be behind their normal stages of development for this time of the year. Most producers are hoping for calm and warm days as well as rainfall to help invigorate crops, hay and pasture.

Crop damage this past week was attributed to dry conditions, wind, flooding, frost and gophers. Some alfalfa is reported to have been damaged by winterkill.

### Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

Strong winds and wet and cool conditions have delayed crop, hay and pasture development in the region. Haying operations are underway with three per cent of the crop cut. Hay quality is rated as 10 per cent excellent, 58 per cent good and 32 per cent fair. Due to excess moisture conditions, some areas remain unseeded while other areas still have un-harvested crops from last year.

Northeast Crop Development - June 26, 2017			
	% Ahead	% Normal	% Behind
Fall Cereals	20	38	42
Spring Cereals	2	44	54
Oilseeds	1	39	60
Pulse Crops	4	42	54

Rainfall across the region ranged from small amounts to 57 mm in the Nipawin area. The Nipawin area holds both the regional and provincial record (360 mm) for the greatest amount of precipitation since April 1.

Topsoil moisture on cropland is rated as 25 per cent surplus, 70 per cent adequate, four per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as 25 per cent surplus, 69 per cent adequate, four per cent short and two per cent very short. Crop District 8A (Nipawin area) reports surplus moisture on 55 per cent of crop land and 45 per cent of hay land and pasture. Crop District 8B (Vonda area) reports only 11 per cent of cropland and 17 per cent of hay land and pasture are short to very short topsoil moisture.

The unfavourable growing conditions have slowed crop development in the region, with 60 per cent of oilseeds, 54 per cent of pulses and spring cereals, and 42 per cent of fall cereals reported to be behind their normal stages of development for this time of the year. Most producers are hoping for calm and warm days to invigorate crops, hay and pasture.

Crop damage this past week was attributed to wet conditions, wind, flooding and frost.

**Northwestern Saskatchewan:**

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Strong winds and wet and cool conditions have delayed crop, hay and pasture development in the region. Haying operations have just begun, with less than one per cent of the crop cut. Hay quality is rated as 33 per cent excellent, 50 per cent good and 17 per cent fair. Due to excess moisture conditions, some areas remain unseeded while other areas still have un-harvested crops from last year.

Northwest Crop Development - June 26, 2017			
	% Ahead	% Normal	% Behind
Fall Cereals	N/A	N/A	N/A
Spring Cereals	0	58	42
Oilseeds	0	57	43
Pulse Crops	0	87	13

Rainfall across the region ranged from small amounts to 40 mm in the Pierceland area. The Pierceland area also holds the record (302 mm) for the greatest amount of precipitation in the region since April 1.

Topsoil moisture on cropland is rated as 16 per cent surplus, 74 per cent adequate and 10 per cent short. Hay land and pasture topsoil moisture is rated as 14 per cent surplus, 71 per cent adequate and 15 per cent short. Crop District 9B (Meadow Lake area) reports surplus moisture on 30 per cent of crop land and 23 per cent of hay land and pasture. Only 10 per cent of cropland and 16 per cent of hay land and pasture is short topsoil moisture.

The unfavourable growing conditions have slowed crop development in the region, with 43 per cent of oilseeds, 13 per cent of pulses and 42 per cent of spring cereals reported to be behind their normal stages of development for this time of year. Most producers are hoping for calm and warm days to help invigorate crops, hay and pasture.

Crop damage this past week was attributed to wet conditions, wind, flooding, frost, cutworms and flea beetles.



## Crop Conditions - June 26, 2017

Provincial							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	9	8	10	13	10	7	1
% good	56	46	60	36	58	57	60
% fair	25	41	25	34	25	30	34
% poor	9	4	4	16	7	5	5
% very poor	1	1	1	1	0	1	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	2	8	0	3	8	6	0
% good	48	52	26	58	57	57	34
% fair	36	29	45	25	25	31	54
% poor	14	10	29	13	9	6	12
% very poor	0	1	0	1	1	0	0
Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	12	21	10	5	8	6	0
% good	55	66	61	35	60	54	52
% fair	26	7	23	39	31	33	46
% poor	5	3	4	18	1	5	2
% very poor	2	3	2	3	0	2	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	4	4	0	2	9	9	0
% good	48	49	22	64	60	52	50
% fair	32	30	71	26	21	34	50
% poor	16	13	7	6	9	4	0
% very poor	0	4	0	2	1	1	0
Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	1	0	2	16	0	2	10
% good	22	27	39	28	43	44	57
% fair	26	68	42	35	44	43	18
% poor	51	5	16	20	13	9	15
% very poor	0	0	1	1	0	2	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	0	1	0	0	1	2	0
% good	11	45	11	5	37	53	29
% fair	55	39	44	8	44	35	57
% poor	34	14	45	87	18	10	14
% very poor	0	1	0	0	0	0	0
East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	4	4	6	0	5	2	3
% good	73	91	61	72	59	58	64
% fair	22	5	29	24	30	35	22
% poor	1	0	4	4	6	5	11
% very poor	0	0	0	0	0	0	0

East-central (continued)							
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	2	4	0	8	4	2	0
% good	59	51	75	64	74	77	85
% fair	34	37	25	24	17	17	15
% poor	5	8	0	4	4	4	0
% very poor	0	0	0	0	1	0	0
West-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	3	19	17	13	17	15	0
% good	43	33	62	60	64	57	63
% fair	53	48	20	25	17	27	35
% poor	1	0	1	2	2	1	2
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	1	19	0	0	19	12	0
% good	59	57	67	62	64	62	75
% fair	39	22	32	37	16	25	25
% poor	1	2	1	1	1	1	0
% very poor	0	0	0	0	0	0	0
Northeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	15	7	17	10	0
% good	0	81	60	70	49	58	71
% fair	0	19	16	20	22	24	21
% poor	0	0	9	3	12	5	8
% very poor	100	0	0	0	0	3	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	0	10	0	13	18	15	N/A
% good	68	51	70	49	50	60	N/A
% fair	29	25	20	37	19	20	N/A
% poor	3	14	9	1	4	5	N/A
% very poor	0	0	1	0	9	0	N/A
Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	N/A	13	N/A	4	4	N/A
% good	0	N/A	62	N/A	86	80	N/A
% fair	0	N/A	16	N/A	8	12	N/A
% poor	0	N/A	6	N/A	2	4	N/A
% very poor	100	N/A	3	N/A	0	0	N/A
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	10	10	N/A	N/A	7	0	N/A
% good	90	61	N/A	N/A	76	94	N/A
% fair	0	22	N/A	N/A	12	6	N/A
% poor	0	6	N/A	N/A	4	0	N/A
% very poor	0	1	N/A	N/A	1	0	N/A

# Crop Districts and Rural Municipalities in Saskatchewan

- Regional Service Office
- Regional Satellite Office
- ▭ Crop District
- ▭ Rural Municipality



Projection: UTM Zone 13 Datum: NAD83




Data Source:  
Crop Districts - Saskatchewan Ministry of Agriculture

Geomatic Services, Ministry of Agriculture June 10, 2014

# Weekly Rainfall

from June 20 to June 26, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



# Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period June 20 to 26, 2017


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	NIL	110	4A	49	White Valley	NIL	57	7A	287	St. Andrews	13	94
	3	Enniskillen	1	94		51	Reno	2.79	54.14		288	Pleasant Valley	9	93
	31	Storthoaks	N/A	10		79	Arlington	2	64		290 A	Kindersley	12.68	100.28
	32	Reciprocity	NIL	124		109	Carmichael	N/A	82		290 B	Kindersley	11	51
	33	Moose Creek	5	116		110	Piapot	7	71.5		290 C	Kindersley	N/A	24
	34	Browning	NIL	99		111	Maple Creek	N/A	67		292	Milton	N/A	113
	61	Antler	N/A	108	4B	139 A	Gull Lake	N/A	52		317 A	Marriott	N/A	N/A
	64	Brock	NIL	116		139 B	Gull Lake	N/A	56.5		317 B	Marriott	12	99
	65	Tecumseh	3	111		142	Enterprise	N/A	N/A		318	Mountain View	14	147
1B	91	Maryfield	NIL	122		169	Pittville	N/A	34		320 A	Oakdale	25	111.5
	94	Hazelwood	5.07	121.52		231	Happyland	7	94		320 B	Oakdale	20	116
	122	Martin	3	98	5A	183	Fertile Belt	4	105		321	Prairiedale	14	125
	123	Silverwood	6	114		211 A	Churchbridge	3	111	7B	347	Biggar	15	108
	124	Kingsley	6	142		211 B	Churchbridge	5	44		350 A	Mariposa	6	120.64
	125 A	Chester	N/A	114		211 C	Churchbridge	N/A	N/A		350 B	Mariposa	5	128
	125 B	Chester	NIL	157		213	Saltcoats	3	97		351	Progress	7	129
	151	Rocanville	3	123		218	Cupar	N/A	N/A		352	Heart's Hill	14	132
	154 A	Elcapo	NIL	112		241	Calder	3	66		377	Glenside	11	111
	154 B	Elcapo	3	107		243	Wallace	NIL	43		378	Rosemount	17	139
	155	Wolseley	2	107.25		244	Orkney	4	68		379	Reford	9	103
2A	67	Weyburn	NIL	58		245 A	Garry	3	65		381	Grass Lake	N/A	64.5
	68	Brokenshell	NIL	64		245 B	Garry	N/A	80		382	Eye Hill	15	127.5
	96	Fillmore	NIL	10.25		245 C	Garry	4	82.5		409 A	Buffalo	5	155
	97	Wellington	NIL	78.5		246 A	Ituna Bon Accord	1	84		409 B	Buffalo	4	132
2B	127 A	Francis	5	121.5		246 B	Ituna Bon Accord	10	96		410	Round Valley	7.59	153.77
	127 B	Francis	5	41.25		247	Kellross	2	85	8A	395	Porcupine	NIL	137
	129	Bratt's Lake	NIL	65		248	Touchwood	8	73		397	Barrier Valley	21.67	125.92
	131 A	Baildon	2	96		271	Cote	2	97		428	Star City	10	103
	131 B	Baildon	12	108		273	Sliding Hills	NIL	83		456	Arborfield	31	182
	156 A	Indian Head	2.49	93		277	Emerald	5	117		457	Connaught	8	166
	156 B	Indian Head	N/A	119	5B	305	Invermay	8	107		486	Moose Range	22	186
	159	Sherwood	4	76		307	Elfros	9	82		487	Nipawin	57	359.5
	160 A	Pense	4	50		308 A	Big Quill	4	61	8B	369	St. Peter	8	73
	160 B	Pense	NIL	50.5		308 B	Big Quill	1	58		370 A	Humboldt	1	74
	161	Moose Jaw	2	63		331	Livingston	9	112		370 B	Humboldt	3	72
	162	Caron	10	68		335	Hazel Dell	10	46		371	Bayne	8	94
	191	Marquis	3	59		336	Sasman	2	78		372	Grant	11.48	106.83
3ASE	38 A	Laurier	2.29	59.69		337	Lakeview	5	124		400	Three Lakes	12	87
	38 B	Laurier	NIL	62		338	Lakeside	7	116		429 A	Flett's Springs	12	109
	39	The Gap	NIL	44		366	Kelvington	17	137		429 B	Flett's Springs	23	150.5
3ASW	10	Happy Valley	NIL	56		367	Ponass Lake	5	131.5		459	Kinistino	10	78
	12	Poplar Valley	NIL	55		368	Spalding	9	101	9AE	460	Birch Hills	30.09	106.65
	40	Bengough	N/A	N/A	6A	190 A	Dufferin	17	118		488	Torch River	10	144
	42	Willow Bunch	NIL	50		190 B	Dufferin	6	112		491	Buckland	10	78.5
	43	Old Post	8	61		190 C	Dufferin	NIL	96		520	Paddockwood	NIL	137
	73 A	Stonehenge	NIL	57.89		190 D	Dufferin	N/A	69	9AW	521	Lakeland	NIL	137
	73 B	Stonehenge	1	74		219 A	Longlaketon	N/A	44		406	Mayfield	8	82
3AN	102	Lake Johnston	2.09	51		219 B	Longlaketon	5	116		435	Redberry	11	159
	103	Sutton	NIL	47		220	McKillop	4	80		436	Douglas	5	118
	132 A	Hillsborough	1.5	43.5		221 A	Sarnia	4.79	94		463	Duck Lake	22	149.62
	132 B	Hillsborough	6	120		221 B	Sarnia	5	93.94		466	Meeting Lake	19	199
	193	Eyebrow	3	55		222	Craik	6	78		467 A	Round Hill	13	158
	17	Val Marie	N/A	20.97		251	Big Arm	NIL	67		467 B	Round Hill	9	163
3BS	18	Lone Tree	5	39		252	Arm River	N/A	51		467 C	Round Hill	18	201
	75	Pinto Creek	1	81		279	Mount Hope	N/A	66.07		494	Canwood	18	157
	76	Auvergne	N/A	61		282	McCraney	N/A	71	9B	438	Battle River	4	142
	77	Wise Creek	2	50		312	Morris	NIL	78.5		440	Hillsdale	10.5	166.5
	78	Grassy Creek	5	68.48		313	Lost River	6	87		442	Manitou Lake	16.98	185.6
	105	Glenbain	NIL	51		339	Leroy	6.59	77.95		498 A	Parkdale	16	185
	106	Whiska Creek	2	92		340	Wolverine	6	96		498 B	Parkdale	N/A	113.5
	107	Lac Pelletier	N/A	44		341	Viscount	2	70		499 A	Mervin	15	186
	108	Bone Creek	NIL	59		343 A	Blucher	11	59		501 A	Frenchman Butte	N/A	201
	138 A	Webb	8	97		223 A	Huron	N/A	0		501 B	Frenchman Butte	12	156
3BN	138 B	Webb	12	75	6B	223 B	Huron	N/A	83		501 C	Frenchman Butte	16	251
	165	Morse	13	105		284 A	Rudy	7	54		502	Britannia	11	208.5
	166	Excelsior	10	73		284 B	Rudy	18	119		561	Loon Lake	29	223
	168 A	Riverside	16	81		284 C	Rudy	14	90.5		588 A	Meadow Lake	39	271
	168 B	Riverside	8	73.23		285	Fertile Valley	9	61		588 B	Meadow Lake	40	302
	226	Victory	4	51		286	Milden	N/A	78		622	Beaver River	N/A	290.7
	228 A	Lacadena	2	109		314	Dundurn	14	72					
	228 B	Lacadena	N/A	6.5		344 A	Corman Park	5	82					
	257	Monet	9.5	88.5		344 B	Corman Park	N/A	103					
						346	Perdue	13	90					
						376	Eagle Creek	13	111					
						403	Rosthern	9	115					

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.


Municipality No: A, B, C and D - more than one reporter

# Cumulative Rainfall

from April 1 to June 26, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.




Data Source:  
Rainfall - Ministry of Agriculture, Crop Report Database  
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture June 28, 2017

# Cropland Topsoil Moisture Conditions


June 26, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

# Hay and Pasture Topsoil Moisture Conditions

## June 26, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.