

Crop Report

Published by the Ministry of Agriculture ISSN 0701 7085 Report number 23, Oct 5, 2017

For the Period September 26 to October 2, 2017

Warm and dry weather earlier in the week allowed many producers to return to the field after several days of rain delays. Eighty-nine per cent of the crop is now in the bin, up from 78 per cent last week. Harvest progress remains ahead of the five year (2012-2016) average of 82 per cent for this time of year. Eight per cent of the crop is swathed or ready to straight-cut.

Harvest is most advanced in the southwestern region, where 98 per cent of the crop is now combined. The southeastern and west-central regions have 92 per cent combined and the east-central region 89 per cent. The northeastern region has 78 per cent combined, while the northwestern region has 69 per cent combined.

Many producers worked long hours over the last week to combine remaining crops prior to the rain. Some crops have been taken off tough and placed into aeration bins.

Ninety-nine per cent of chickpeas, 98 per cent of mustard, 97 per cent of durum, 93 per cent of canaryseed, 92 per cent of spring wheat and barley, 84 per cent of canola, 81 per cent of oats, 64 per cent of flax and 46 per cent of soybeans have now been combined.

Fourteen per cent of canola is swathed or ready to straight-cut.

Harvest Progress in SK Per cent Combined All Crops								
Oct 2/17	89							
5 year avg. (2012-2016)	82							
Oct 3/16	80							
Oct 5/15	84							
Oct 6/14	72							
Sept 30/13	79							
Oct 1/12	97							
10 year avg. (2007-2016)	83							

Producers in many areas of the province
welcomed the heavy rain and snow to
help replenish the topsoil; however,
warm and dry weather is now needed to
dry crops sufficiently for producers to
finish harvest. Rainfall this past week
ranged from small amounts to 92 mm in
the Kenaston area. Many central and
northeastern areas reported well over 50
mm of rainfall, while areas in the south
reported snow.

One	year	ago
-----	------	-----

Eighty per cent of the crop was now in the bin. Heavy rain and snow halted harvest progress for many areas of the province and fields remained saturated. Spring wheat quality was below long-term averages.

Follow the 2017 Crop Report on Twitter @SKAgriculture

Saskatchewan Harvest October 2, 2017							
Per cent cor							
Winter wheat	100						
Fall rye*	100						
Spring wheat	92						
Durum	97						
Oats**	81						
Barley	92						
Canaryseed	93						
Flax	64						
Canola	84						
Mustard	98						
Soybeans	46						
Lentils	99						
Peas	99						
Chickpeas 99							
*includes four per cent 'other'							
**includes three per							
cent 'other'							

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.saskatchewan.ca/crop-report.





Topsoil moisture conditions have greatly improved with the recent moisture, but more will be needed to replenish the subsoil moisture for next spring. Across the province, topsoil moisture on cropland is rated as five per cent surplus, 39 per cent adequate, 43 per cent short and 13 per cent very short. Hay land and pasture topsoil moisture is rated as two per cent surplus, 34 per cent adequate, 39 per cent short and 25 per cent very short.

Yield estimates remain about average overall, although they vary greatly across the province depending on time of seeding and moisture received throughout the season. Some eastern and northern areas are reporting higher-than-average yields, thanks to timely rainfall. Yields in much of the southern and central regions were greatly affected by the extended period of hot and dry conditions this summer. Many crops did not fill properly as they ran out of moisture much earlier than normal. Some cereal crops have reduced protein content and lighter bushel weights. Spring wheat grades are above the 10 year average and are being reported as 74 per cent 1 CW, 23 per cent 2 CW and three per cent 3 CW.

The majority of crop damage this past week was due to localized flooding, frost, strong winds, lack of moisture and wildlife such as geese and deer.

Producers are busy combining, completing fall field work, moving cattle and hauling bales.

SaskPower received five reports last week of farm machinery coming in contact with electrical equipment, bringing the total for September to 28. Producers are urged to be especially careful when using equipment around power lines. Safety information is available at www.saskpower.com/safety.

	Saskatchewan Harvest by Crop District October 2, 2017 % combined											
1A	1A 92 3BS 97 7A 92											
1B	95	3BN	98	7B	96							
2A	89	4A	98	8A	75							
2B	94	4B	99	8B	84							
3ASE	97	5A	87	9AE	72							
3ASW	97	5B	81	9AW	69							
3AN	96	6A	99	9B	69							
		6B	90									

	Provincial Estimated Crop Yields - October 2, 2017											
	Winter wheat	Fall rye	HRSW	Other wheat*	Durum	Oat	Barley	Canaryseed				
Southeast	41	41	38	41	33	68	58	1159				
Southwest	27	24	28	40	28	50	44	975				
East Central	50	45	44	47	41	83	65	1303				
West Central	40	35	42	42	39	64	59	1156				
Northeast	70	36	49	65	50	105	68	1234				
Northwest	57	55	46	54	-	87	61	-				
Provincial	42	38	42	48	31	86	60	1172				
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea					
Southeast	18	28	704	20	30	1270	900					
Southwest	13	22	676	15	23	1142	887					
East Central	25	35	1173	27	40	1375	1100					
West Central	24	35	1145	23	39	1686	-					
Northeast	27	39	-	30	45	1500	-					
Northwest	25	38	-	-	44	1324	-					
Provincial	20	34	786	21	32	1334	910					

^{* &#}x27;Other wheat' includes all wheat classes other than Hard Red Spring Wheat

Southeastern Saskatchewan:

- Crop District 1 Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE Radville and Lake Alma areas

Thanks to a stretch of warm and dry weather, producers were able to return to the field. Ninety-two per cent of the crop is now in the bin, up from 88 per cent last week and well ahead of the five-year (2012-2016) average for this time of year of 84 per cent. While the majority of producers have completed harvest, others are waiting on crops such as flax and soybeans to dry.

Rainfall in the region was welcomed last week to help replenish the topsoil moisture; however, significant rain is still needed to replenish the subsoil for next year. Rainfall over the weekend ranged from 1 mm in the Broadview and Maryfield areas to 30 mm in the Moose Jaw area. Many areas reported receiving at least half an inch of moisture. The Alameda area has reported the most precipitation (267 mm) in the region since April 1.

Topsoil moisture conditions have significantly improved with the recent moisture. Topsoil moisture on cropland is rated as 40 per cent adequate, 42 per cent short and 18 per cent

^{**} Crop yield predictions at this point in time. Please keep in mind these are regional averages, and yields can vary greatly across an area.

^{***} canaryseed, mustard, lentil and chickpea in lbs/ac. All other crops in bu/ac.

very short. Hay land and pasture topsoil moisture is rated as 30 per cent adequate, 47 per cent short and 23 per cent very short. Although field conditions have improved in much of the region, Crop District 3ASE continues to report that 100 per cent of cropland, hay land and pasture are short to very short topsoil moisture at this time.

Yields are estimated to be about average in the region but vary greatly. Those areas that received timely rainfall are reporting slightly above-average yields, while other areas that missed the rains are reporting well-below-average yields. Spring wheat grades at this point in time are being reported as 71 per cent 1 CW, 24 per cent 2 CW, five per cent 3 CW and one per cent CW Feed. There are reports of lighter-than-normal bushel weights and reduced protein content in cereals.

Most crop damage this past week was due to strong winds, frost and lack of moisture. Post-harvest herbicide applications continue on those fields that have active weed growth. Livestock producers are concerned about feed shortages and some have already begun shipping cattle.

Producers are busy finishing harvest, working fields and cleaning corrals.

Southwestern Saskatchewan:

- Crop District 3ASW Coronach, Assiniboia and Ogema areas
- Crop District 3AN Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 Consul, Maple Creek and Leader areas

Those producers with crops remaining in the field were able to get back in last week and continue harvesting. Ninety-eight per cent of the crop is now in the bin, up from 92 per cent last week and well ahead of the five-year (2012-2016) average of 88 per cent for this time of year. Producers are waiting for crops such as flax and soybeans to dry.

Heavy rain and snow were welcomed in the region to help replenish topsoil moisture. Additional moisture will be needed to replenish the subsoil for next year. Rain and snow this past week ranged from trace amounts to 78 mm in the Tyner area. Many western areas of the region reported well over two inches of moisture. The Tyner area has reported the most precipitation (226 mm) in the region since April 1.

Cropland topsoil moisture conditions continue to improve with the recent moisture and, for the first time in over four months, surplus topsoil moisture has been reported. Topsoil moisture on cropland is rated as one per cent surplus, 15 per cent adequate, 60 per cent short and 24 per cent very short. Hay land and pasture topsoil moisture is rated as 14 per cent adequate, 35 per cent short and 51 per cent very short.

Yield estimates vary greatly across the region but are below average overall. There are some areas reporting higher-than-expected yields on crops such as lentils. The extended period of hot and dry conditions greatly affected production in the region. Spring wheat grades at this point in time are being reported as 75 per cent 1 CW, 22 per cent 2 CW and three per cent 3 CW.

Crop damage this week is attributed to frost, strong winds and lack of moisture. Post-harvest herbicide applications continue on those fields that have active weed growth. Livestock producers are concerned about feed shortages and some have already begun shipping cattle.

Producers are busy completing harvest, working fields and moving cattle.

East-Central Saskatchewan:

- Crop District 5 Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A Lumsden, Craik, Watrous and Clavet areas

Producers in the region put in long hours this past week to harvest the remaining crop before the rain. Eighty-nine per cent of the crop is now in the bin, up from 77 per cent last week and well ahead of the five-year (2012-2016) average of 74 per cent for this time of year. Some producers have been taking the crop off tough and placing it into aeration bins. Heavy rain earlier this week halted harvest, and most producers need a week or more of good weather in order to complete harvest.

Large amounts of rain fell on the region, helping to replenish the topsoil. Rainfall this past week ranged from trace amounts to 92 mm in the Kenaston area. There are reports of rain exceeding 100 mm in some areas of the region. The Pelly area has reported the most precipitation (347 mm) in the region since April 1.

Topsoil moisture conditions have improved with the rain. Topsoil moisture on cropland is rated as 12 per cent surplus, 48 per cent adequate, 34 per cent short and six per cent very short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 47 per cent adequate, 38 per cent short and nine per cent very short. Crop District 6A is reporting that 44 per cent of the cropland and 52 per cent of the hay land and pasture are short to very short topsoil moisture at this time, a dramatic improvement over last week. Yield estimates at this time vary greatly across the region. Overall yields are average to above-average, depending on moisture received through the summer. Spring wheat grades at this point in time are being reported as 81 per cent 1 CW, 18 per cent 2 CW and one per cent 3 CW.

Crop damage this week is attributed to localized flooding, frost, strong winds and wildlife such as geese and deer. Post-harvest herbicide applications continue on those fields that have active weed growth.

Producers are busy combining, baling straw, moving cattle and working fields.

West-Central Saskatchewan:

- Crop District 6B Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A Rosetown, Kindersley, Eston, Major
- Crop District 7B Kerrobert, Macklin, Wilkie and Biggar areas

Thanks to some warm and dry days earlier in the week, producers in the region were able to get back into the field. Ninety-two per cent of the crop is now in the bin, up from 81 per

cent last week and well ahead of the five-year (2012-2016) average of 84 per cent for this time of year. Some producers are taking off crops tough and placing them in aeration bins. Heavy rain late in the week halted harvest, and most producers need a week or more of good weather in order to complete harvest.

Rain and snow this past week ranged from trace amounts to 69 mm in the Rosetown area. While most areas received at least 35 mm, some areas reported localized downpours of well over 75 mm. The Tramping Lake area has reported receiving the most precipitation (298 mm) in the region since April 1.

Topsoil moisture has drastically improved with the recent rainfall; however, more rain will be needed to replenish the subsoil. Topsoil moisture on cropland is rated as two per cent surplus, 31 per cent adequate, 50 per cent short and 17 per cent very short. Hay land and pasture topsoil moisture is rated as two per cent surplus, 25 per cent adequate, 51 per cent short and 22 per cent very short.

Yield estimates vary greatly across the region, but are about average overall. There are some areas reporting higher-than-expected yields on crops such as lentils. The extended period of hot and dry conditions affected production in much of the region. Spring wheat grades at this point in time are being reported as 77 per cent 1 CW, 22 per cent 2 CW and one per cent 3 CW.

The majority of crop damage this past week was due to frost, strong winds, lack of moisture and wildlife such as deer and geese. Post-harvest herbicide applications continue on those fields that have active weed growth.

Producers are busy completing harvest, working fields, hauling bales and moving cattle.

Northeastern Saskatchewan:

- Crop District 8 Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE Prince Albert, Choiceland and Paddockwood areas

A quarter of the region's crops were harvested last week, thanks to some warm and dry weather. Many producers worked long hours to combine remaining crops before the weekend rain; some crop is coming off tough and being placed in aeration bins.

Seventy-eight per cent of the crop is now in the bin, up from 53 per cent last but slightly below the five-year (2012-2016) average of 80 per cent for this time of year. Several weeks of warm and dry weather are needed in order for producers to finish.

Heavy rainfall was received in much of the region, ranging from 3 mm in the Garrick and Birch Hills areas to 78 mm in the Porcupine Plain area. Many areas reported receiving at least 25 mm of rain. The Porcupine Plain area has reported the most precipitation (344 mm) in the region since April 1.

Across the region, topsoil moisture conditions on cropland are rated as 16 per cent surplus, 59 per cent adequate and 25 per cent short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 62 per cent adequate and 33 per cent short.

Yield estimates vary throughout the region, but range from about average to well above average in the region. Timely moisture throughout the season positively affected production for most producers. Spring wheat grades at this point in time are being reported as 70 per cent 1 CW, 25 per cent 2 CW, four per cent 3 CW and one per cent CW Feed.

Crop damage this past week was due to strong winds, localized flooding and wildlife such as elk, deer and waterfowl. Post-harvest herbicide applications continue on those fields that have active weed growth.

Producers are busy combining, moving cattle and working fields.

Northwestern Saskatchewan:

- Crop District 9AW Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Producers put in long hours this past week to combine remaining crop before the weekend rains. Sixty-nine per cent of the crop is now in the bin, up significantly from 48 per cent last. The five-year (2012-2016) average for this time of year is 84 per cent combined. While some producers have wrapped up, many more will need several weeks of warm and dry weather in order to do so. Some crops are coming off tough and being placed into aeration bins.

While most of the region only received small amounts of rain, the Neilburg area reported 42 mm. The Meadow Lake area has reported the most precipitation (484 mm) in both the region and the province since April 1.

Cropland topsoil moisture conditions are rated as 65 per cent adequate, 29 per cent short and six per cent very short. Hay land and pasture topsoil moisture is rated as 54 per cent adequate, 34 per cent short and 12 per cent very short.

Yield estimates vary throughout the region, but range from about average to well above average in the region. Timely moisture throughout the season positively affected production for most producers. Spring wheat grades at this point in time are being reported as 66 per cent 1 CW, 30 per cent 2 CW and four per cent 3 CW.

The majority of crop damage this past week is attributed to frost, strong winds and wildlife such as geese and deer. Some crops have been downgraded due to sprouting and bleaching caused by the recent rains. Post-harvest herbicide applications continue on those fields that have active weed growth.

Producers are busy combining, moving cattle, working fields and hauling bales.


Saskatchewan Harvest Progress - October 2, 2017

*Other - crop that will not be harvested due to weather, insect or disease damage or will be greenfeed

Other - crop that will not be harvested due to weather, insect of disease damage of will be greenieed											
Winter Wheat	% Standing	% in swath	% ready to straight combine	% combined							
southeast	0	0	0	100							
southwest	0	0	0	100							
east central	0	0	0	100							
west central	0	0	0	100							
northeast	0	0	0	100							
northwest	0	0	0	100							
provincial	0	0	0	100							
Fall Rye	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)						
southeast	0	0	otraight combine	97	(greenieed/snage)						
southwest	0	0	0	95	5						
east central	0	0	0	100	0						
west central	0	0	0	100	0						
northeast	0	0	0	100	0						
northwest	0	0	0	100	0						
provincial	0	0	0	96	4						
			% ready to	% combined	-						
Spring Wheat	% Standing	% in swath	straight combine								
southeast	0	0	2	98							
southwest	0	0	2	98							
east central	2	2	2	94							
west central	1	1	4	94							
northeast	2	1	7	90							
northwest	8	5	14	73							
provincial	2	2	4	92							
Durum	% Standing	% in swath	% ready to straight combine	% combined							
southeast	1	0	1	98							
southwest	0	0	1	99							
east central	0	0	0	100							
west central	8	2	4	86							
northeast	0	0	0	100							
northwest	N/A	N/A	N/A	N/A							
provincial	2	0	1	97							
Barley	% Standing	% in swath	% ready to straight combine	% combined							
southeast	0	0	0	100							
southwest	0	0	1	99							
east central	1	2	2	95							
west central	2	2	3	93							
northeast	3	2	2	93							
northwest	16	11	8	65							
provincial	3	2	3	92							
Oats	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)						
southeast	0	0	1	97	2						
southwest	0	0	8	80	12						
east central	6	5	7	80	2						
west central	6	5	2	71	16						
northeast	4	2	14	79	1						
northwest	15	14	14	55	2						
provincial	6	4	9	78	3						
Canaryseed	% Standing	% in swath	% ready to straight combine	% combined							
southeast	2	1	0	97							
southwest	8	0	0	92							
east central	0	0	0	100							
west central	0	0	2	98							
northeast	21	2	18	59							
northwest	N/A	N/A	N/A	N/A							
provincial	3	1	3	93							


Flax	% Standing	% in swath	% ready to straight combine	% combined	
southeast	25	5	6	64	
southwest	18	0	12	70	
east central	16	7	5	72	
west central	30	6	5	59	
northeast	47	21	10	22	
northwest	28	0	0	72	
provincial	24	6	6	64	
Canola	% Standing	% in swath	% ready to straight combine	% combined	
southeast	3	2	2	93	
southwest	2	0	2	96	
east central	2	10	2	86	
west central	1	7	1	91	
northeast	1	23	5	71	
northwest	3	22	10	65	
provincial	2	11	3	84	
Mustard	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	14	86	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	0	0	2	98	
Soybeans	% Standing	% in swath	% ready to straight combine	% combined	
southeast	36	0	18	46	
southwest	36	0	20	44	
east central	26	0	26	48	
west central	20	0	6	74	
northeast	61	0	30	9	
northwest	N/A	N/A	N/A	N/A	
provincial	35	0	19	46	
Field Peas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	0	0	1	99	
provincial	0	0	1	99	
Lentils	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	1	99	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0 00 00 00 00 00	0	1 % ready to	99	
Chickpeas	% Standing	% in swath	straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	1	99	
east central	0	0	0	100	
west central	N/A	N/A	N/A	N/A	
			/ .		
northeast northwest	N/A N/A	N/A N/A	N/A N/A	N/A N/A	

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

from September 26 to October 2, 2017



Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm


for the period Sept 26 to Oct 2, 2017

1A 1B	2 3 31 32 33 34 61 64 65 91 94 122 123 124 125 A	Mount Pleasant Enniskillen Storthoaks Reciprocity Moose Creek Browning Antler Brock Tecumseh Maryfield Hazelwood Martin	4 3 N/A 18 12 N/A 2	220 218 10 258 267	4A	49 51	White Valley	N/A	93	7A	287	St. Andrews	N/A	
1B	31 32 33 34 61 64 65 91 94 122 123 124 125 A	Storthoaks Reciprocity Moose Creek Browning Antler Brock Tecumseh Maryfield Hazelwood	N/A 18 12 N/A 2	10 258		51								167
1B	32 33 34 61 64 65 91 94 122 123 124 125 A	Reciprocity Moose Creek Browning Antler Brock Tecumseh Maryfield Hazelwood	18 12 N/A 2	258			Reno	7.6	97.1		288	Pleasant Valley	69	222
1B	33 34 61 64 65 91 94 122 123 124 125 A	Moose Creek Browning Antler Brock Tecumseh Maryfield Hazelwood	12 N/A 2			79	Arlington	15	106		290 A	Kindersley	52.7	207.8
1B	34 61 64 65 91 94 122 123 124 125 A	Browning Antler Brock Tecumseh Maryfield Hazelwood	N/A 2	267		109	Carmichael	N/A	95		290 B	Kindersley	NIL	87.4
1B	61 64 65 91 94 122 123 124 125 A	Antler Brock Tecumseh Maryfield Hazelwood	2			110	Piapot	N/A	79		290 C	Kindersley	N/A	24
1B	64 65 91 94 122 123 124 125 A	Brock Tecumseh Maryfield Hazelwood		204		111	Maple Creek	45	141		292	Milton	N/A	196
1B	65 91 94 122 123 124 125 A	Tecumseh Maryfield Hazelwood		302	4B	139 A	Gull Lake	48	151		317 A	Marriott	N/A	N/A
1B	91 94 122 123 124 125 A	Maryfield Hazelwood		201		139 B	Gull Lake	N/A	69.5		317 B	Marriott	7	194
18	94 122 123 124 125 A	Hazelwood	9	229		169	Pittville	N/A	119		318	Mountain View	8	239
	122 123 124 125 A		1	232		231	Happyland	41	206		320 A	Oakdale	37	232
	123 124 125 A		N/A	209.2	5A	183 211 A	Fertile Belt	10 8	191		320 B	Oakdale Prairiedale	N/A 58	202 258
	124 125 A	Silverwood	15 5	223 195		211 A 211 B	Churchbridge Churchbridge	N/A	272 44	7B	321 347	Biggar	3	179
	125 A	Kingsley	1	209		211 6	Saltcoats	6	189	/ b	350 A	Mariposa	35	298.1
		Chester	24	180		241	Calder	20	187		350 A	Mariposa	18	207
			16	233		243	Wallace	NIL	153		351	Progress	N/A	231
	151	Rocanville	25	246		244	Orkney	7	172		352	Heart's Hill	55	237
		Elcapo	N/A	143		245 A	Garry	N/A	157		377	Glenside	2	218
		Elcapo	N/A	123		245 B	Garry	19	156		378	Rosemount	4	292
	155	Wolseley	9.5	177		245 C	Garry	N/A	117		379	Reford	14	198
2A	67	Weyburn	15	135		246 A	Ituna Bon Accord	25	167		381	Grass Lake	48	216
	68	Brokenshell	11	118		246 B	Ituna Bon Accord	16.1	195.6		382	Eye Hill	27	266.9
	96	Fillmore	14	33		247	Kellross	23	138		409 A	Buffalo	6	219
	97	Wellington	14	126		248	Touchwood	45	163		409 B	Buffalo	5	247
2B		Francis	10	176	5B	271	Cote	16	236		410	Round Valley	12.5	293.9
	127 B	Francis	17.5	83.8		273	Sliding Hills	10	227	8A	395	Porcupine	78	344
	129	Bratt's Lake	10.5	118.5		277	Emerald	24	264		397	Barrier Valley	34.4	304.2
	131 A	Baildon	19	153		305	Invermay	26	273		426	Bjorkdale	50	95
		Baildon	30	198		307	Elfros	40	244		428	Star City	N/A	162
		Indian Head	17.3	170.2		308 A	Big Quill	59	185		456	Arborfield	43	331
		Indian Head	20	211		308 B	Big Quill	75	207		457	Connaught	42	248
	159	Sherwood	12	123.1		331	Livingston	40	347		486	Moose Range	12	303
	160 A		N/A	88		335	Hazel Dell	N/A	113		487	Nipawin	N/A	553.5
		Pense	25	113.2		336	Sasman	N/A	211	8B	369	St. Peter	23	166
	161	Moose Jaw	26	144		337	Lakeview	N/A	215		370 A	Humboldt	37	207
	162	Caron	19	155		338	Lakeside	N/A	196		370 B	Humboldt	26	183
ACE	191	Marquis	7.3	91		366 367	Kelvington	38	343		371	Bayne	18	216
BASE		Laurier Laurier	5	112.8 114		368	Ponass Lake Spalding	55 35	322.5 311		372 400	Grant Three Lakes	12.6 27	233.8 232
	39	The Gap	11	96	6A	190 A	Dufferin	16	175		429 A	Flett's Springs	18	180
ASW	10	Happy Valley	14	114	0.7	190 A	Dufferin	13	165		429 B	Flett's Springs	23	225
AOVV	12	Poplar Valley	8	101		190 C	Dufferin	24	149		459	Kinistino	15	147
	42	Willow Bunch	8	105		190 D	Dufferin	20	106		460	Birch Hills	3.2	148.9
	43	Old Post	NIL	102		219 A	Longlaketon	N/A	95	9AE	488	Torch River	3	241
		Stonehenge	26.7	114.1		219 B	Longlaketon	N/A	145	0712	491	Buckland	5	170.5
	73 B	Stonehenge	24	177.5		220	McKillop	31	161		520	Paddockwood	5	196.5
	74	Wood River	26.7	45		221 A	Sarnia	N/A	159.9		521	Lakeland	5	196.5
3AN	102	Lake Johnston	19.6	125.3		221 B	Sarnia	N/A	127.1	9AW	406	Mayfield	NIL	212
	103	Sutton	26	106		222	Craik	54	173		435	Redberry	N/A	277
	132 A	Hillsborough	32	133		251	Big Arm	65	185.5		436	Douglas	NIL	207
		Hillsborough	30	226		252	Arm River	52	192		463	Duck Lake	9	267.6
	193	Eyebrow	45	148		279	Mount Hope	51.9	203		466	Meeting Lake	2	319
3BS	17	Val Marie	8.5	44.1		282	McCraney	92	241		467 A	Round Hill	NIL	273
	18	Lone Tree	NIL	66		312	Morris	80	238.5		467 B	Round Hill	N/A	297
	75	Pinto Creek	25	135		313	Lost River	N/A	173		467 C	Round Hill	2	399
	76	Auvergne	31	122		339	Leroy	34.8	215.2		493	Shellbrook	NIL	68.8
	77	Wise Creek	32	125		340	Wolverine	32	273		494	Canwood	3	272
	78	Grassy Creek	3	95.5		341	Viscount	N/A	123		497	Medstead	2	92.5
	105	Glenbain	25	131	0.5	343	Blucher	20	203	9B	438	Battle River	12	247
	106	Whiska Creek	13	144	6B	223 A		68	190		440	Hillsdale	32.5	349
	107	Lac Pelletier	34	116.5		223 B		50	147		442	Manitou Lake	41.7	376.2
ODA!	108	Bone Creek	N/A	108		284 A	Rudy	35	214		498 A	Parkdale	N/A	354
3BN	138 A		47	210		284 B		28	224.5		498 B	Parkdale Mervin	N/A	288.5
	138 B		52 N/A	169		284 C		32	192		499 501 A		18	448.1
	165	Morse Excelsion	N/A	77		285	Fertile Valley	25	164		501 A	Frenchman Butte	2	416
	166	Excelsior Riverside	70 64	176		286 314	Milden Dundurn	45	232		501 B	Frenchman Butte Frenchman Butte	N/A	338
		Riverside Riverside	52	181 174.5		314 344	Corman Park	32 NIL	185 219		501 C 502	Britannia	N/A N/A	404 274.5
	226	Victory	N/A	51		344 346	Perdue	7	155		502 561	Loon Lake	2 N/A	435
		Lacadena	78	226		346 376	Eagle Creek	NIL	208		561 588 A	Meadow Lake	3	435
		Lacadena	N/A	6.5		403	Rosthern	NIL	244		588 B	Meadow Lake	5	484
	257	Monet	77	208		403	MOSITION	INIL	244		622	Beaver River	N/A	563.4
	201	INIOLICE		200							022	Deaver Mivel	IN/A	505.4

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.


Cumulative Rainfall

from April 1 to October 2, 2017




Cropland Topsoil Moisture Conditions

October 2, 2017




Hay and Pasture Topsoil Moisture Conditions

October 2, 2017



Minimum Temperature

from September 26 to October 2, 2017



Maximum Temperature

from September 26 to October 2, 2017

